

H2 MA 2019 JC2 Prelim (Paper 1 and Paper 2)

Filename: Change SCHOOL to your school name, e.g. NYJC

Paper 1

Select topic from the dropdown list. If the question consists of multiple topics, choose 1 topic.

QN	TOPIC (H2) Paper 1	ANSWERS (Exclude graphs and text answers)
1	Equations & Inequalities	72 units
2	Sigma Notation & MOD	(i) $\ln \frac{1+x}{1+x^{N+1}}$ (ii) $\ln(1+x)$
3	Maclaurin & Binomial Series	$p = \sqrt{3}, q = 1, r = \frac{1}{2\sqrt{3}}$
4	Graphs & Transformations	(i) $D = (-\infty, -1] \cup [1, \infty)$ (ii) $y = \pm \frac{x+5}{6}$
5	Vectors	(i) $\overrightarrow{OM} = \frac{1}{3}(3\mathbf{a} + \lambda\mathbf{b}), \overrightarrow{ON} = \frac{4}{7}\lambda\mathbf{b}$ (ii) $k = \frac{1}{3}$
6	Differentiation & Applications	(i) $\frac{dy}{dx} = \frac{-1}{25 \cos t}$ (ii) $\frac{225}{4\sqrt{2}}$ (iii) $y = 0$
7	Integration & Applications	(a)(i) $-\sin \frac{x}{2} \left(e^{\cos \frac{x}{2}} \right)$ (ii) $-2 \cos \frac{x}{2} \left(e^{\cos \frac{x}{2}} \right) + 2e^{\cos \frac{x}{2}} + C$ (b) $-\ln 1 - e^x + x + C$
8	Functions	(ii) greatest value of k is a . (iii) $f^{-1} : x \mapsto a - 2\sqrt{a^2 - x^2}, x \in \mathbb{R}, 0 \leq x \leq a$ (iv) $R_{\text{gf}} = \left[\frac{\sqrt{3}}{2}a, a \right]$
9	APGP	(i) $d = 268.0$ (ii) 1339.9 cm (iii) red

		(iv) yellow
10	Complex Numbers	(a) $u = \frac{a}{4+a^2} - \frac{6+2a^2}{4+a^2}i, v = \frac{2}{4+a^2} - \frac{a}{4+a^2}i$ (b)(i) $k = 12$ (iii) $b(x+12)(x + \frac{1+\sqrt{1+4b^2}}{2b}i)(x + \frac{1-\sqrt{1+4b^2}}{2b}i)$
11	Differentiation & Applications	(a) -0.916 cm/min (b) -2.29 cm/min (c) 4293.510 cm ³
12	Vectors	(ii) 11.1° (iv) $(0.547, -0.237, 0.00325)$ (v) 117 m
13	H2 Prelim P1 Q13 Topic	
14	H2 Prelim P1 Q14 Topic	

Paper 2

Select topic from the dropdown list. If the question consists of multiple topics, choose 1 topic.

QN	TOPIC (H2) Paper 2	ANSWERS (<u>Exclude graphs and text answers</u>)
1	Equations & Inequalities	(i) $x \leq -\frac{a}{2}$ or $x = a$ or $x \geq \frac{5a}{2}$ (ii) $x \leq -a$ or $x = \frac{a}{2}$ or $x \geq 2a$
2	Maclaurin & Binomial Series	(ii) $y = 1 + x^2 - \frac{3}{2}x^3 + \dots$ (iii) 0.985 (iv) $\frac{1}{e^{\sin x} \sqrt{1-2x}} \approx 1 + x^2 + \frac{3}{2}x^3$
3	Complex Numbers	(i) $\frac{a}{2}; -\frac{\pi}{3}$ (ii) $-\frac{5\pi}{12}; a \cos \frac{\pi}{12}$ (iii) 6
4	Differential Equations	(a) (ii) $P = 2\sqrt{1+3e^{-2t}}$ (iii) The population of the bugs will decrease and approach 2000 in the long run. (b) $N = 4t \ln t + t$
5	Normal Distribution	(i) 7.6
6	Correlation & Regression	(ii) $(69, 53)$ (iii) Since $ r = 0.96785$ for Model B is nearest to 1, Model B is the most accurate model (iv) 62 marks (v) unreasonable as 120 is not within data range

		where $48 \leq x \leq 84$
7	Normal Distribution	(i) a “6-inch” loaf more likely to be less than 6 inches (ii) $\frac{1}{2}$ (iii) 0.855 (iv) 0.446
8	PnC & Probability	(a) (i) 3360 (ii) 1200 (iii) 240 (b) (i) $\frac{13}{14}$ (ii) 0.644
9	Hypothesis Testing	(a) (i) 949.84 ; 1,73 (ii) $H_0: \mu = 950$ $H_1: \mu < 950$ (iii) 0.135 (iv) $\{\alpha \in \mathbb{R} : 13.5 \leq \alpha \leq 100\}$ (b) $12.5 \leq k \leq 104.8$ (c) There is no need for the floor supervisor to assume the volume of shower gel follow a normal distribution as the sample sizes in both part (a) and (b) are large, the sample mean volume of shower gel can be approximated to follow a normal distribution by Central Limit Theorem.
10	DRV	(i) \$2.50; \$3.75 (ii) Option A is a “sure win” option where the player would definitely gain a positive amount in all cases, whereas option B has a risk of losing money in some cases. (iv) $A \sim N\left(125, \frac{125}{2}\right)$; $B \sim N\left(\frac{375}{2}, \frac{22175}{8}\right)$
11	H2 Prelim P2 Q11 Topic	
12	H2 Prelim P2 Q12 Topic	
13	H2 Prelim P2 Q13 Topic	
14	H2 Prelim P2 Q14 Topic	

Qn	Solutions	C
1	$D(x) = \frac{40320}{g(x)}$ $\Rightarrow g(x) = \frac{40320}{D(x)}$ $ax^2 + bx + c = \frac{40320}{D(x)}$ <p>Given $5^2a + 5b + c = \frac{40320}{384} = 105$ -- (1)</p> $8^2a + 8b + c = \frac{40320}{224} = 180$ -- (2) $10^2a + 10b + c = \frac{40320}{168} = 240$ -- (3) <p>Using GC, $a = 1, b = 12, c = 20$</p> <p>When $x = 18,$</p> $D(18) = \frac{40320}{18^2 + 12(18) + 20} = 72 \text{ units}$	<p>G w A g(5^2 of 5^2</p> <p>A de in pc</p>
2i	<p><u>Method 1: (method of differences)</u></p> $\sum_{n=1}^N p_n = \sum_{n=1}^N \ln \frac{1+x^n}{1+x^{n+1}}$ $= \sum_{n=1}^N (\ln(1+x^n) - \ln(1+x^{n+1}))$ $= \ln(1+x) - \ln(1+x^2)$ $+ \ln(1+x^2) - \ln(1+x^3)$ $+ \ln(1+x^3) - \ln(1+x^4)$ \vdots $+ \ln(1+x^N) - \ln(1+x^{N+1})$ $= \ln(1+x) - \ln(1+x^{N+1})$ $= \ln \frac{1+x}{1+x^{N+1}}$	<p>TI dc dc su di cl th Sc en lin co se</p> <p>A fir ln of</p>


Method 2: (using property of logarithm)

$$\begin{aligned} \sum_{n=1}^N p_n &= \sum_{n=1}^N \ln \frac{1+x^n}{1+x^{n+1}} \\ &= \ln \frac{1+x}{1+x^2} + \ln \frac{1+x^2}{1+x^3} + \ln \frac{1+x^3}{1+x^4} + \dots + \ln \frac{1+x^N}{1+x^{N+1}} \\ &= \ln \frac{1+x}{1+x^2} \cdot \frac{1+x^2}{1+x^3} \cdot \frac{1+x^3}{1+x^4} \cdot \dots \cdot \frac{1+x^N}{1+x^{N+1}} \\ &= \ln \frac{1+x}{1+x^{N+1}} \end{aligned}$$

2ii Since $-1 < x < 1$, as $N \rightarrow \infty$, $x^{N+1} \rightarrow 0$.

$$\begin{aligned} \therefore \sum_{n=1}^{\infty} p_n &= \ln \frac{1+x}{1+0} \\ &= \ln(1+x) \end{aligned}$$

3



By Cosine Rule,

$$\begin{aligned} \cos\left(\frac{\pi}{2} - x\right) 2^2 &= 1^2 + (BC)^2 - 2(BC)\cos\left(\frac{\pi}{2} - x\right) \\ 4 &= 1 + (BC)^2 - 2(BC)\sin x \end{aligned}$$

$$(BC)^2 - 2(BC)\sin x - 3 = 0$$

$$BC = \frac{2\sin x \pm \sqrt{4\sin^2 x + 12}}{2}$$

$$= \sin x \pm \sqrt{\sin^2 x + 3}$$

$$\approx x \pm \sqrt{x^2 + 3}$$

$$= x + \sqrt{x^2 + 3}, \text{ since } BC > 0$$


$$= x + \sqrt{3} \left[1 + \frac{1}{3}x^2 + \dots \right]$$

$$\approx \sqrt{3} + x + \frac{x^2}{2\sqrt{3}}$$

$$p = \sqrt{3}, q = 1, r = \frac{1}{2\sqrt{3}}$$

$$p = \sqrt{3}, q = 1, r = \frac{1}{2\sqrt{3}}$$

Alternative Solution



By sine rule,

$$\frac{BC}{\sin\left(\frac{\pi}{2} + x - y\right)} = \frac{2}{\sin\left(\frac{\pi}{2} - x\right)} = \frac{1}{\sin y}$$

$$\Rightarrow \frac{BC}{\cos(y - x)} = \frac{2}{\cos x} = \frac{1}{\sin y}$$

$$\Rightarrow BC = \frac{2 \cos(y - x)}{\cos x}$$

$$\Rightarrow BC = \frac{2(\cos y \cos x + \sin y \sin x)}{\cos x}$$


$$\Rightarrow BC = \frac{2(\cos y \cos x + \sin y \sin x)}{\cos x}$$

$$\Rightarrow BC = \frac{2\left(\frac{\sqrt{4 - \cos^2 x}}{2} \cos x + \frac{\cos x}{2} \sin x\right)}{\cos x}$$

$$\Rightarrow BC = \frac{\sqrt{4 - \cos^2 x} \cos x + \cos x \sin x}{\cos x}$$

$$\Rightarrow BC = \sqrt{4 - \cos^2 x} + \sin x$$

$$\Rightarrow BC = \sqrt{3 + \sin^2 x} + \sin x$$



- 4i** Sketch the graph using GC, remember to set window appropriately.


OR
KIASU ExamPaper

Notice that it is the upper half of a hyperbola.

$$y = \sqrt{\frac{(x+5)^2}{36} - 1}$$


$$\frac{(x+5)^2}{6^2} - y^2 = 1$$

We have $D = (-\infty, -1] \cup [1, \infty)$.


<p>ii</p>	<p>Let $y = \sqrt{\frac{(x+5)^2}{36} - 1}$</p> $\frac{(x+5)^2}{6^2} - y^2 = 1$ <p>To find asymptotes:</p> $y^2 = \frac{(x+5)^2}{6^2}$ $y = \pm \frac{x+5}{6}$
<p>iii</p>	 <p>The graph shows a hyperbola $y = f(x)$ on a Cartesian coordinate system. The vertices are marked at $(-11, 0)$ and $(1, 0)$. The origin is labeled O. Two dashed lines represent the asymptotes, with equations $y = \frac{1}{6}x + \frac{5}{6}$ and $y = -\frac{1}{6}x - \frac{5}{6}$.</p>

iv


Using GC to help if students is not sure



$$y = \frac{1}{f(x)}$$



$$y = f'(x)$$



5i

$$\overline{OC} = \lambda \mathbf{b}$$

$$\overline{OD} = \overline{OA} + \overline{OC} = \mathbf{a} + \lambda \mathbf{b}$$

$$\overline{ON} = \frac{4}{7} \lambda \mathbf{b}$$

$$\overline{OM} = \frac{\overline{OD} + 2\overline{OA}}{3}$$

$$= \frac{\mathbf{a} + \lambda \mathbf{b} + 2\mathbf{a}}{3}$$

$$= \frac{1}{3}(3\mathbf{a} + \lambda \mathbf{b})$$

ii

Area of triangle $OMD = \frac{1}{2} |\overline{OM} \times \overline{OD}|$

$$= \frac{1}{2} \left| \frac{1}{3} (\lambda \mathbf{b} + 3\mathbf{a}) \times (\lambda \mathbf{b} + \mathbf{a}) \right|$$

$$= \frac{1}{6} \left| \lambda^2 \mathbf{b} \times \mathbf{b} + \lambda \mathbf{b} \times \mathbf{a} + 3\lambda \mathbf{a} \times \mathbf{b} + 3\mathbf{a} \times \mathbf{a} \right|$$

$$= \frac{1}{6} \left| -\lambda \mathbf{a} \times \mathbf{b} + 3\lambda \mathbf{a} \times \mathbf{b} \right| \quad (\because \mathbf{b} \times \mathbf{b} = \mathbf{0} = \mathbf{a} \times \mathbf{a})$$

$$= \frac{1}{3} \lambda |\mathbf{a} \times \mathbf{b}|$$

iii

$|\mathbf{p} \cdot \mathbf{a}|$ is the length of projection of \overline{OA} on \overline{OD}

6i $\frac{dx}{dt} = 50 \sin t \cos t$, $\frac{dy}{dt} = -2 \sin t$
 $\frac{dy}{dx} = \frac{-1}{25 \cos t}$

ii When $t = \frac{\pi}{4}$, $\frac{dy}{dx} = \frac{-\sqrt{2}}{25}$, $x = \frac{25}{2}$, $y = \sqrt{2}$

$$y = -\frac{\sqrt{2}}{25}x + c$$

$$c = \sqrt{2} + \frac{1}{\sqrt{2}} = \frac{3}{\sqrt{2}}$$

Equation of tangent: $y = -\frac{\sqrt{2}}{25}x + \frac{3}{\sqrt{2}}$

When $x = 0$, $y = \frac{3}{\sqrt{2}}$

When $y = 0$, $x = \frac{75}{2}$

Area of triangle $OBO = \frac{1}{2} \left(\frac{75}{2} \right) \left(\frac{3}{\sqrt{2}} \right) = \frac{75}{4} \left(\frac{3}{\sqrt{2}} \right) = \frac{225}{4\sqrt{2}}$


KIASU
 Exam Apps
 Islandwide Delivery | Whatsapp Only 88660031

iii	$y = 0$
7ai	<p>Let $y = 2e^{\cos \frac{x}{2}}$</p> $\frac{dy}{dx} = -\frac{1}{2} \sin \frac{x}{2} \left(2e^{\cos \frac{x}{2}} \right)$ $= -\sin \frac{x}{2} \left(e^{\cos \frac{x}{2}} \right)$
aii	$\int \frac{1}{2} \sin x \left(e^{\cos \frac{x}{2}} \right) dx$ $= \int \frac{1}{2} \left(2 \sin \frac{x}{2} \cos \frac{x}{2} \right) \left(e^{\cos \frac{x}{2}} \right) dx$ $= -\int \left(\cos \frac{x}{2} \right) \left(-\sin \frac{x}{2} e^{\cos \frac{x}{2}} \right) dx$ $= \left(-\cos \frac{x}{2} \right) \left(2e^{\cos \frac{x}{2}} \right) - \int \left(\frac{1}{2} \sin \frac{x}{2} \right) \left(2e^{\cos \frac{x}{2}} \right) dx$ $= -2 \cos \frac{x}{2} \left(e^{\cos \frac{x}{2}} \right) + 2e^{\cos \frac{x}{2}} + C$
b	$\int \frac{1}{1-e^x} dx$ <p>Let $u = 1 - e^x \Rightarrow \frac{du}{dx} = -e^x = u - 1 \Rightarrow \frac{dx}{du} = \frac{1}{u-1}$</p> $= \int \frac{1}{u(u-1)} du$ $= \int \left(-\frac{1}{u} + \frac{1}{u-1} \right) du$ $= -\ln u + \ln u-1 + C$ <p style="text-align: right;">www.KiasuExamPaper.com 312</p>

$$= -\ln|1 - e^x| + \ln|-e^x| + C$$

$$= -\ln|1 - e^x| + x + C$$

8i



ii

For f^{-1} to exist, f must be one-one. i.e. every horizontal line $y = h$, $h \in \mathbb{R}$ can only cut the graph of $y = f(x)$ at most once. Hence, the greatest value of k is a .

iii

$$\text{Let } y = \sqrt{a^2 - \frac{(x-a)^2}{4}}$$

$$\Rightarrow y^2 = a^2 - \frac{(x-a)^2}{4}$$

$$\Rightarrow (x-a)^2 = 4(a^2 - y^2)$$

$$\Rightarrow x = a \pm 2\sqrt{a^2 - y^2}$$

$$\Rightarrow x = a - 2\sqrt{a^2 - y^2} \quad (\because -a \leq x \leq a)$$


$$f: x \mapsto a - 2\sqrt{a^2 - x^2}, x \in \mathbb{R}, 0 \leq x \leq a$$

wiv

$$g(x) = f\left(\frac{3}{2}x\right) \text{ for } x \in \mathbb{R}, -\frac{2}{3}a \leq x \leq 2a.$$

Since $R_f = [0, a] \subseteq \left[-\frac{2}{3}a, 2a\right] = D_g$, the composite


function gf exists.



$$D_{gf} = D_f \xrightarrow{f} R_f = [0, a] \xrightarrow{g} \left[\frac{\sqrt{3}}{2} a, a \right] = R_{gf}$$

$$R_{gf} = \left[\frac{\sqrt{3}}{2} a, a \right]$$

9i



$$d\left(\frac{4}{5}\right)^7 = 56.2$$

$$\therefore d = 267.9824829 = 268.0 \text{ (1 d.p.)}$$

ii

Since $|r| = \left|\frac{4}{5}\right| < 1$, sum to infinity exists.

Hence maximum theoretical length of wire


$$= \frac{d}{1-r}$$

$$= \frac{267.9824829}{1 - \frac{4}{5}}$$

$$= 1339.912415$$

$$= 1339.9 \text{ cm (1 d.p.)}$$

iii



Let S_n be distance from LED 1 to LED $n+1$.

Method 1: (using GC table)

Using sum of GP, $S_n = \frac{267.9824829[1 - (\frac{4}{5})^n]}{1 - \frac{4}{5}}$

Using GC,

n	S_n	$ S_n - 1290 $
14	1281.0	9.0
15	1292.8	2.8

\therefore closest LED to 1290 is when $n = 15$.

\therefore required LED is LED $(15+1) =$ LED 16.

Hence colour of LED 16 is red.

Method 2: (using algebraic manipulation)

Let $S_n = \frac{267.9824829[1 - (\frac{4}{5})^n]}{1 - \frac{4}{5}} = 1290$

$$1 - (\frac{4}{5})^n = 0.9627494943$$

$$n \ln(\frac{4}{5}) = \ln 0.0372505057$$

$$\therefore n = 14.74427443$$

Using GC,

n	S_n	$ S_n - 1290 $
14	1281.0	9.0
15	1292.8	2.8

\therefore closest LED to 1290 is when $n = 15$.


\therefore required LED is LED $(15+1) =$ LED 16.

Hence colour of LED 16 is red.

KIASU
ExamPaper

Islandwide Delivery | Whatsapp Only 88660031

iv



$$\text{Let } u_n = d \left(\frac{4}{5}\right)^{n-1} = 267.9824829 \left(\frac{4}{5}\right)^{n-1} \geq 1$$

$$\left(\frac{4}{5}\right)^{n-1} \geq \frac{1}{267.9824829}$$

$$\ln\left(\frac{4}{5}\right)^{n-1} \geq \ln \frac{1}{267.9824829}$$

$$(n-1) \ln\left(\frac{4}{5}\right) \geq \ln \frac{1}{267.9824829}$$

$$n-1 \leq 25.05526859$$

$$\therefore n \leq 26.05526859$$

Since largest integer $n = 26$, last LED is LED $(26+1) =$
LED 27

Hence colour of last LED 27 is yellow.

10a $v + iu = 2$ ----(1)

$$av - 2u = 3i$$
 ----(2)

$$(1) \times a - (2):$$

$$iau + 2u = 2a - 3i$$

$$u = \frac{2a - 3i}{2 + ia} \times \frac{2 - ia}{2 - ia}$$

$$u = \frac{4a - 6i - 2a^2i - 3a}{4 + a^2}$$

$$u = \frac{a - (6 + 2a^2)i}{4 + a^2}$$

$$= \frac{a}{4 + a^2} - \frac{6 + 2a^2}{4 + a^2}i$$

Sub u into (1)

$$v = 2 - iu$$

$$= 2 - i\left(\frac{a}{4 + a^2} - \frac{6 + 2a^2}{4 + a^2}i\right)$$

$$= \frac{2(4 + a^2) - a^2 + 6 + 2a^2}{4 + a^2} + \frac{a}{4 + a^2}i$$

Islandwide Delivery | Whatsapp Only 88660031

bi	<p>Sub $x = -k$ into</p> $bx^3 + (12b + i)x^2 + (12i + b)x + 12b = 0$ <p>We have:</p> $-bk^3 + (12b + i)k^2 - (12i + b)k + 12b = 0$ $-bk^3 + 12bk^2 - bk + 12b = 0 \quad \text{--- (1)}$ <p>and $k^2 - 12k = 0 \quad \text{--- (2)}$</p> <p>(2) implies</p> $k(k - 12) = 0$ $k = 0, 12$ <p>Reject $k = 0$ as $b \neq 0$.</p>
bii	$bx^2 + ix + b = 0$ $x = \frac{-i \pm \sqrt{-1 - 4b^2}}{2b}$ $x = \frac{-1 \pm \sqrt{1 + 4b^2}}{2b} i$ <p>Hence roots are purely imaginary</p> <div style="text-align: center;">  <p>KIASU ExamPaper</p> <p><small>Islandwide Delivery Whatsapp Only 88660031</small></p> </div>
biii	<p>Hence</p> $bx^3 + (12b + i)x^2 + (12i + b)x + 12b = (x + 12)(bx^2 + cx + b)$ <p>By compare coeff of x, $c = -317$, using (i):</p>

$$bx^3 + (12b+i)x^2 + (12i+b)x + 12b = (x+12)(bx^2 + ix + b)$$

$$= b(x+12)\left(x + \frac{1 + \sqrt{1+4b^2}}{2b}i\right)\left(x + \frac{1 - \sqrt{1+4b^2}}{2b}i\right)$$

11 Height of liquid in the cylinder = $x - 4$

a

$$V = \pi(2.5)^2(x-4) + \frac{1}{3}\pi(2.5)^2(4)$$

$$= 6.25\pi(x-4) + \frac{25}{3}\pi$$

$$\frac{dV}{dx} = 6.25\pi \quad \text{and} \quad \frac{dV}{dt} = -18$$

$$\frac{dx}{dt} = \frac{dx}{dV} \times \frac{dV}{dt}$$

$$= \frac{-18}{6.25\pi}$$

$$= -0.916 \text{ cm/min}$$

b

$$\frac{r}{x} = \frac{2.5}{4} \Rightarrow x = 1.6r$$

When $x = 2$, $r = \frac{5}{8}$

KIASU ExamPaper 

Islandwide Telegram Whatsapp Only 88660031

$$V_c = \frac{1}{3}\pi r^2 x$$

$$= \frac{1}{3}\pi r^2(1.6r)$$

$$= \frac{1.6}{3}\pi r^3$$

$$\frac{dV}{dr} = 1.6\pi r^2$$

$$\frac{dr}{dt} = \frac{dr}{dV} \times \frac{dV}{dt}$$

$$= \frac{1}{1.6\pi \left(\frac{5}{4}\right)^2} \times (-18)$$

$$= -2.29 \text{ cm/min}$$

c

$$\text{Vol of bowl} = \pi \int_{-10}^0 225 \left(1 - \frac{y^2}{100}\right) dy$$

Vol. of empty space

$$= \pi \int_{-10}^0 225 \left(1 - \frac{y^2}{100}\right) dy - \pi (2.5)^2 (20) - \frac{1}{3} \pi (2.5)^2 (4)$$

$$= 4293.510 \text{ cm}^3 \text{ (3 d.p.)}$$

A

B

12i

$$\text{Equation of line through } MD : \mathbf{r} = \begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 7 \\ -1 \end{pmatrix}, \lambda \in \mathbb{R}$$

$$\begin{pmatrix} 0.8 \\ 0.75 \\ 0.3 \end{pmatrix} = \begin{pmatrix} 0.4 \\ 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 0.4 \\ 1.65 \\ 0.3 \end{pmatrix}$$

$$\text{Equation of line through } ND : \mathbf{r} = \begin{pmatrix} 0.4 \\ -0.9 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 0.4 \\ 1.65 \\ 0.3 \end{pmatrix}, \mu \in \mathbb{R}$$

Since lines through MD and ND intersect at D ,

St
the
lin
r :

\bar{M}

W
In

(
(

	$\begin{cases} 0.8 + 2\lambda = 0.4 + 0.4\mu \\ 0.6 + 7\lambda = -0.9 + 1.65\mu \\ -\lambda = 0.3\mu \end{cases}$ <p>Solving using GC,</p> $\lambda = -\frac{3}{25}, \mu = \frac{2}{5}.$ <p>Substitute $\lambda = -1.2$ into equation of line through MD,</p> $\begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} - \frac{3}{25} \begin{pmatrix} 2 \\ 7 \\ -1 \end{pmatrix} = \begin{pmatrix} 0.56 \\ -0.24 \\ 0.12 \end{pmatrix}$ <p>\therefore coordinates of D is $(0.56, -0.24, 0.12)$. (shown)</p>	Ma ans or sho cur
ii	<p>Equation of horizontal ground: $\mathbf{r} \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 0$</p> <p>Required angle</p> $= 90^\circ - \cos^{-1} \frac{\begin{vmatrix} \begin{pmatrix} 0.56 \\ -0.24 \\ 0.12 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \\ \sqrt{0.56^2 + (-0.24)^2 + 0.12^2} \sqrt{1} \end{vmatrix}}$ $= 11.14233567^\circ$ $= 11.1^\circ \text{ (1 d.p.)}$	Ma for for Ma no: ho: So equ wr is 1
iii	<p>Given $T(0, 0, 0.07)$.</p> $\vec{NM} = \begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} - \begin{pmatrix} 0.4 \\ -0.9 \\ 0 \end{pmatrix} = \begin{pmatrix} 0.4 \\ 1.5 \\ 0 \end{pmatrix}$ $\vec{TM} = \begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} - \begin{pmatrix} 0 \\ 0 \\ 0.07 \end{pmatrix} = \begin{pmatrix} 0.8 \\ 0.6 \\ -0.07 \end{pmatrix}$ $\begin{pmatrix} 0.4 \\ 1.5 \\ 0 \end{pmatrix} \times \begin{pmatrix} 0.8 \\ 0.6 \\ -0.07 \end{pmatrix} = \begin{pmatrix} 0.105 \\ 0.028 \\ -0.96 \end{pmatrix} = 0.01 \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix}$ <p>\therefore equation of p:</p> $\mathbf{r} \cdot \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix} = \begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix} = -6.72 \text{ (Shown)}$	Ma 0.0 $\begin{pmatrix} - \\ \\ \\ \end{pmatrix}$ Stu equ wr pre

iv Equation of line perpendicular to p passing through D is

$$\mathbf{r} = \begin{pmatrix} 0.56 \\ -0.24 \\ 0.12 \end{pmatrix} + t \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix}, t \in \mathbb{R}.$$

Substitute equation of line into p ,

$$\begin{pmatrix} 0.56 - 10.5t \\ -0.24 + 2.8t \\ 0.12 - 96t \end{pmatrix} \cdot \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix} = -6.72$$

$$\therefore t = 0.00121618712$$

Hence

$$\begin{pmatrix} 0.56 \\ -0.24 \\ 0.12 \end{pmatrix} + 0.00121618712 \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix} = \begin{pmatrix} 0.5472300352 \\ -0.2365946761 \\ 0.00324603648 \end{pmatrix}$$

$$= \begin{pmatrix} 0.547 \\ -0.237 \\ 0.00325 \end{pmatrix}$$

\therefore coordinates of G is $(0.547, -0.237, 0.00325)$.

v Method 1: [Hence using $(0.547, -0.237, 0.00325)$]

Required distance DG

$$= \sqrt{(0.56 - 0.54723)^2 + (-0.24 - (-0.23659))^2 + (0.12 - 0.003246)^2}$$

$$= 0.1174998$$

$$= 0.117 \text{ (3 s.f.)}$$

\therefore required distance = 117 m

Method 2: [Otherwise using dot product]

$$\vec{MD} = \begin{pmatrix} 0.56 \\ -0.24 \\ 0.12 \end{pmatrix} - \begin{pmatrix} 0.8 \\ 0.6 \\ 0 \end{pmatrix} = \begin{pmatrix} -0.24 \\ -0.84 \\ 0.12 \end{pmatrix}$$

Required distance

$$= \left| \begin{pmatrix} -0.24 \\ -0.84 \\ 0.12 \end{pmatrix} \cdot \frac{\begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix}}{\sqrt{(-10.5)^2 + 2.8^2 + (-96)^2}} \right|$$

$$= 0.1174996006$$

$$= 0.117 \text{ (3 s.f.)}$$

\therefore required distance = 117 m

Method 3


$$|\overrightarrow{DG}| = \beta \begin{pmatrix} -10.5 \\ 2.8 \\ -96 \end{pmatrix}$$

$$= 0.00121618712\sqrt{9334.09}$$

$$= 0.1174996006$$


$$= 0.117 \text{ (3 s.f.)}$$

\therefore required distance = 117 m


Qn	Solutions	Comments
1		<p>In general part.</p> <p>Common</p> <ol style="list-style-type: none"> 1. Fo in 2. Th st of 3. Sc nu gr
ii	<p>To find points of intersection, we solve</p> $y = 3ax - 3a^2 $ $y = 2(x - a)^2$ <p>i.e. $2(x - a)^2 = 3ax - 3a^2$</p> $2(x - a)^2 = 3ax - 3a^2 \text{ ----- (1)}$ $\Rightarrow 2x^2 - 7ax + 5a^2 = 0$ $\Rightarrow (2x - 5a)(x - a) = 0$ $\Rightarrow x = \frac{5}{2}a \text{ or } x = a$ <p>Or $2(x - a)^2 = -(3ax - 3a^2) \text{ ----- (2)}$</p> $\Rightarrow 2x^2 - ax - a^2 = 0$ $\Rightarrow (2x + a)(x - a) = 0$ $\Rightarrow x = -\frac{1}{2}a \text{ or } x = a$ <p>For $2(x - a)^2 \geq 3ax - 3a^2$,</p> $x \leq -\frac{a}{2} \text{ or } x = a \text{ or } x \geq \frac{5a}{2}$	<p>A lot of s previous</p> <p>Majority that $x = a$</p> <p>The easie intercepts based on final conc</p> <p>A lot of s method to directly w manipulat</p> <p>Common</p> <ol style="list-style-type: none"> 1. a m M lo 2. Sc w in fa Th (

HWA

1ii	<p>Replace x with $x + \frac{a}{2}$,</p> $2\left(x + \frac{a}{2} - a\right)^2 \geq \left 3a\left(x + \frac{a}{2}\right) - 3a^2\right $ $2\left(x - \frac{a}{2}\right)^2 \geq \left 3ax - \frac{3a^2}{2}\right $ $\therefore x + \frac{a}{2} \leq -\frac{a}{2} \text{ or } x + \frac{a}{2} = a \text{ or } x + \frac{a}{2} \geq \frac{5a}{2}$ $\Rightarrow x \leq -a \text{ or } x = \frac{a}{2} \text{ or } x \geq 2a$	A lot replac
2i	$y = \frac{e^{\sin x}}{\sqrt{1+2x}}$ $y\sqrt{1+2x} = e^{\sin x}$ $\sqrt{1+2x} \frac{dy}{dx} + \frac{1}{2} \frac{2}{\sqrt{1+2x}} y = \cos x e^{\sin x}$ $\frac{dy}{dx} + \frac{y}{1+2x} = \frac{e^{\sin x}}{\sqrt{1+2x}} \cos x$ $= y \cos x$ $\frac{1}{y} \frac{dy}{dx} + \frac{1}{1+2x} = \cos x$ <div style="text-align: center;">  <p>Alternative Exam Paper Islandwide Delivery Whatsapp: 98866688</p> </div> $\frac{\sqrt{1+2x} \left(\frac{dy}{dx}\right) + (1+2x)^{-\frac{1}{2}}}{y\sqrt{1+2x}} = \cos x$ $\frac{1}{y} \frac{dy}{dx} + \frac{1}{1+2x} = \cos x \text{ (Shown)}$	This c Stude detail quest credit

<p>ii</p>	<p>Differentiating again w.r.t. x,</p> $\frac{1}{y} \frac{d^2 y}{dx^2} - \frac{1}{y^2} \left(\frac{dy}{dx} \right)^2 - \frac{2}{(1+2x)^2} = -\sin x$ $\frac{1}{y} \frac{d^3 y}{dx^3} - \frac{1}{y^2} \frac{dy}{dx} \frac{d^2 y}{dx^2} - \frac{2}{y^2} \frac{dy}{dx} \frac{d^2 y}{dx^2} + \frac{2}{y^3} \left(\frac{dy}{dx} \right)^3 + \frac{8}{(1+2x)^3} = -\cos x$ $\frac{1}{y} \frac{d^3 y}{dx^3} - \frac{3}{y^2} \frac{dy}{dx} \frac{d^2 y}{dx^2} + \frac{2}{y^3} \left(\frac{dy}{dx} \right)^3 + \frac{8}{(1+2x)^3} = -\cos x$ <p>When $x = 0, y = 1, \frac{dy}{dx} = 0, \frac{d^2 y}{dx^2} = 2, \frac{d^3 y}{dx^3} = -9$.</p> <p>The Maclaurin's series for y is</p> $y = 1 + \frac{2}{2!} x^2 - \frac{9}{3!} x^3 + \dots$ $= 1 + x^2 - \frac{3}{2} x^3 + \dots$	<p>The n quest differ $\frac{1}{y} \frac{dy}{dx}$ $\frac{dy}{dx}$ a $\frac{d^2 y}{dx^2}$ expre differ $\frac{d}{dx} \left(\frac{1}{y} \right)$</p>
<p>iii</p>	$\int_0^1 \frac{e^{\sin x}}{\sqrt{1+2x}} dx \approx \int_0^1 \left(1 + x^2 - \frac{3}{2} x^3 \right) dx$ $= \frac{23}{24}$ $= 0.985 \text{ (3 s.f.)}$	<p>A lot they c integr evalu</p>
<p>iii</p>	$\int_0^1 \frac{e^{\sin x}}{\sqrt{1+2x}} dx = 0.058$  <p>KIASU ExamPaper Islandwide Delivery Whatsapp Only 88660031</p> <p>The graphs of $y = \frac{e^{\sin x}}{\sqrt{1+2x}}$ and $y = 1 + x^2 - \frac{3}{2} x^3$ differ significantly near to $x = 1$. Hence the approximation is not good.</p>	<p>1. Ma the M powe powe appro answe expla good.</p> <p>2. Ma that x witho affect</p> <p>3. So range range for ap valid</p>

		not sr less tl good
iv	<p>Replace x with $-x$</p> $y = \frac{e^{\sin(-x)}}{\sqrt{1-2x}} = \frac{e^{-\sin x}}{\sqrt{1-2x}} = \frac{1}{e^{\sin x} \sqrt{1-2x}}$ $\frac{1}{e^{\sin x} \sqrt{1-2x}} \approx 1 + x^2 + \frac{3}{2}x^3$	Many series conne (ii).
3i	$ p = \left \frac{a}{1 + \sqrt{3}i} \right = \frac{a}{2}$ $\arg(p) = \arg\left(\frac{a}{1 + \sqrt{3}i}\right)$ $= \arg(a) - \arg(1 + \sqrt{3}i)$ $= 0 - \frac{\pi}{3}$ $= -\frac{\pi}{3}$	<ul style="list-style-type: none"> • In: arg, mu, p, mi • A 4th • A
ii	<p>The shape is a rhombus.</p>	<ul style="list-style-type: none"> • Al fo ve p sh • Sc di • A qu tri
ii	<p>KIASU Exam Paper Islandwide Delivery Whatsapp Only 88060031</p> $\frac{\pi}{2} + \frac{\pi}{3} = \frac{\pi}{12}$ $\arg(p + q) = -\left(\frac{\pi}{3} + \frac{\pi}{12}\right) = -\frac{5\pi}{12}$	<ul style="list-style-type: none"> • Sc ar on ro • Sc ar an



$$|p + q| = 2 \times OS = 2 \times \frac{a}{2} \cos \frac{\pi}{12} = a \cos \frac{\pi}{12}$$


iii For $(p + q)^n$ to be purely imaginary,
 $(p + q)^n = (a \cos \frac{\pi}{12})^n [\cos(-\frac{5n\pi}{12}) + i \sin(-\frac{5n\pi}{12})]$
 $\cos(-\frac{5n\pi}{12}) = 0$
 Hence the smallest positive integer n is 6, as
 $\cos(-\frac{5\pi}{2}) = 0.$

- M
- fo
- Sc
- ar

- Sc
- su
- a f

<p>4ai</p>	$\frac{dP}{dt} = \frac{a}{P} - bP, \text{ where } a, b \text{ are constants}$ <p>When $P = 2$,</p> $\frac{dP}{dt} = \frac{a}{2} - 2b = 0 \Rightarrow a = 4b$ $\frac{dP}{dt} = \frac{4b}{P} - bP$ $= b \left(\frac{4}{P} - P \right)$ $= k \left(\frac{4}{P} - P \right), \text{ where } k = b$
<p>4aii</p>	<p>When $P = 4$, $\frac{dP}{dt} = -3$.</p> $\frac{dP}{dt} = k \left(\frac{4}{4} - 4 \right) = -3$ $\Rightarrow k = 1$ $\frac{dP}{dt} = \frac{4}{P} - P$ $= \frac{4 - P^2}{P}$ $\frac{dt}{dP} = \frac{P}{4 - P^2}$ $= -\frac{1}{2} \frac{-2P}{4 - P^2}$ $t = -\frac{1}{2} \ln 4 - P^2 + C$ $-2t = \ln 4 - P^2 - 2C$ $\ln 4 - P^2 = 2C - 2t$ $4 - P^2 = Ae^{-2t}, \text{ where } A = \pm e^{2C}$ <p>When $t = 0, P = 4$.</p> $A = -12$ $4 - P^2 = -12e^{-2t}$ $P^2 = 4 + 12e^{-2t}$ $P = 2\sqrt{1 + 3e^{-2t}}$

4a
iii



The population of the bugs will decrease and approach 2000 in the long run.

4b

Method 1:

$$u = \frac{N}{t}$$

$$\frac{du}{dt} = \frac{t \frac{dN}{dt} - N}{t^2}$$

$$t \frac{du}{dt} = \frac{dN}{dt} - \frac{N}{t}$$

Substitute into $\frac{dN}{dt} = 4 + \frac{N}{t}$:

$$t \frac{du}{dt} = 4$$

$$\frac{du}{dt} = \frac{4}{t}$$

$$u = 4 \ln t + C$$

$$\frac{N}{t} = 4 \ln t + C$$

When $t = 1$, $N = 1 \Rightarrow C = 1$.

$$N = 4t \ln t + t$$

Method 2: 
Islandwide Delivery Whatsapp Only 88660031

t

$$N = ut$$

$$\frac{dN}{dt} = \frac{du}{dt} t + u$$

Substitute into $\frac{dN}{dt} = 4 + \frac{N}{t}$:


	$\frac{du}{dt}t + u = 4 + u$ $\frac{du}{dt}t = 4$ $\frac{du}{dt} = \frac{4}{t}$ $u = 4 \ln t + C$ $\frac{N}{t} = 4 \ln t + C$ <p>When $t = 1$, $N = 1 \Rightarrow C = 1$.</p> $N = 4t \ln t + t$	
5 (i)	<p>Let X denote the daily rainfall in mm. $X \sim N(\mu, \sigma^2)$</p> <p>In 7 days, $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{7}\right)$</p> $P(X > 9.8) = 0.1$ $P\left(Z > \frac{9.8 - \mu}{\sigma}\right) = 0.1$ $\frac{9.8 - \mu}{\sigma} = 1.2815516$ $9.8 = \mu + 1.2815516\sigma \quad \text{--- (1)}$ $P(\bar{X} > 8.2) = 0.1$ $P\left(Z > \frac{8.2 - \mu}{\sigma/\sqrt{7}}\right) = 0.1$ $\frac{8.2 - \mu}{\sigma/\sqrt{7}} = 1.2815516$ $8.2 = \mu + 0.4843810\sigma \quad \text{--- (2)}$ <p>Solving (1) and (2), $\mu = 7.22780, \sigma = 2.00710$ $\therefore \sigma = 2.01$ (2dp) (shown)</p>	So μ pa
(ii)	<p>In 30 days, $\bar{X} \sim N\left(7.22780, \frac{2.01^2}{30}\right)$</p> $P(\bar{X} > k) \geq 0.1$ $k \leq 7.6981(5sf)$ $\therefore \max k = 7.6(1dp)$	N al A th So ar

Note: change in inequality sign because for area to be greater than or equal to 0.1, the value of k can be at 7.6974 or to its left, i.e. less than 7.6974 (see diagram below).



Possible actual boundary 7.6974
to get area larger than 0.1

6
(i)



(ii)

(x_7, y_7) is (\bar{x}, \bar{y})

Using G.C,

$(\bar{x}, \bar{y}) = (69, 53)$

OR

$(x_7, y_7) =$

$$\left(\frac{80 + 84 + 70 + 74 + 58 + 48}{6}, \frac{44 + 40 + 49 + 45 + 58 + 82}{6} \right)$$

KIASU
Exam Paper
6

(iii)

Model A: $y = a + bx^2$ r -value = -0.92386

Model B: $y = \ln a + b \ln x$ r -value = -0.96785

Model C: $y = a + b\sqrt{x}$ r -value = -0.95855

Since $|r|$ for Model B is nearest to 1, Model B is the most accurate model

(iv)	$y = \ln a + b \ln x$ <p>By GC,</p> $\ln a = 350.11$ $b = -70.469$ $y = 350.11 - 70.469 \ln x$ <p>When $x = 60$, $y = 62$ marks.</p> <p>The student is estimated to score 62 marks.</p>	N us of m A w A ec
(v)	$4 \times 30 = 120$ <p>unreasonable as 120 is not within data range where $48 \leq x \leq 84$.</p>	So ho Th th ar th m
7 (i)	<p>Let X and Y denote the length (in inches) of a “footlong” and a “6-inch” loaf respectively.</p> $X \sim N(12.2, 0.2^2), Y \sim N(6.1, (0.1\sqrt{2})^2)$ $P(Y < 6) = 0.239750 = 0.240(3sf)$ $P(X < 12) = 0.158655 = 0.159(3sf) < P(Y < 6)$ <p>\therefore a “6-inch” loaf more likely to be less than 6 inches.</p>	A cc va th 6, ga It pr St fu as qu
(ii)	$E(Y_1 + Y_2 - X) = 2(6.1) - 12.2 = 0$ $Var(Y_1 + Y_2 - X) = 2(0.1\sqrt{2})^2 + 0.2^2 = 0.08$ $Y_1 + Y_2 - X \sim N(0, 0.08)$ $P(Y_1 + Y_2 > X) = P(Y_1 + Y_2 - X > 0) = \frac{1}{2}$	It in M th th So w as ar

(iii)	<p>Let A denote the number of “6-inch” sandwiches less than 6-inches in length in a week. $A \sim B(3, 0.239750)$ $P(A \leq 1) = 0.855122 = 0.855(3sf)$</p> <p><u>Alternative</u> required probability = $P(Y < 6)[P(Y > 6)]^2 \times 3 + [P(Y > 6)]^3$ $= 0.23975(1 - 0.23975)^2 \times 3 + (1 - 0.23975)^3$ $= 0.855(3sf)$</p>	St va w or di of sa “1
(iv)	<p>Let B and C denote the number of “6-inch” sandwiches less than 6-inches in length in a 4-week and 3-week period respectively. $B \sim B(12, 0.239750)$ and $C \sim B(9, 0.239750)$</p> $P(A = 1 B > 4) = \frac{P(A = 1)P(C > 3)}{P(B > 4)}$ $= \frac{P(A = 1)[1 - P(C \leq 3)]}{1 - P(B \leq 4)}$ $= \frac{0.41571 \times 0.14710}{0.13721}$ $= 0.445674$ $= 0.446(3sf)$	Th ch fr So cc So de nu 4 th P = So 2 up P = = =
8 (a)(i)	<p>Number of arrangements = $\frac{8!}{3!2!} = 3360$</p>	So re
8 (a)(ii)	<p>Number of arrangements = $\frac{5!}{2!} \times {}^6C_3 = 1200$</p>	M cc 2
8 (a)(iii)	<p>Arrange 3 vowels (into a group) $\frac{3!}{2!} = 3$</p>	M es er

Arrange 5 consonants (for slotting)

$$\frac{5!}{3!} = 20$$

Since the ends must be consonants, only the middle 4 slots can be used for the group to slot. (ie. 4C_1)

$$\text{Number of arrangements} = 3 \times 20 \times 4 = 240$$

OR

Case 1: First and Last 'L'

$$4! \times \frac{3!}{2!} = 72$$

Case 2: First and last 'L' and non-L

$$2 \times 2 \times \frac{4!}{2!} \times \frac{3!}{2!} = 144$$

Case 3: First and last non-L

$$2 \times \frac{4!}{3!} \times \frac{3!}{2!} = 24$$

Total = 240 ways

(b)(i)

$P(1V) + P(2V) + P(3V)$

$$= \frac{{}^5C_3 {}^3C_1 + {}^5C_2 {}^3C_2 + {}^5C_1 {}^3C_3}{{}^8C_4}$$

$$= \frac{65}{70} = \frac{13}{14}$$

Probability

$= 1 - P(\text{all consonants})$

$$= 1 - \left(\frac{5}{8} \times \frac{4}{7} \times \frac{3}{6} \times \frac{2}{5} \right) = \frac{13}{14}$$

OR

$1 - P(\text{no vowels})$

$$= 1 - \frac{{}^5C_4}{{}^8C_4}$$

$$= 1 - \frac{5}{70}$$

$$= \frac{13}{14}$$


OR

$P(1 \text{ vowel}) + P(2 \text{ vowels}) + P(3 \text{ vowels})$

$$\begin{aligned} &= \frac{3 \times 5 \times 4 \times 3}{8 \times 7 \times 6 \times 5} \times \frac{4!}{3!} + \frac{3 \times 2 \times 5 \times 4}{8 \times 7 \times 6 \times 5} \times \frac{4!}{2!2!} + \frac{3 \times 2 \times 1 \times 5}{8 \times 7 \times 6 \times 5} \times \frac{4!}{3!} \\ &= \frac{13}{14} \end{aligned}$$

8

(b)(ii)



Probability

$= P(VCC) + P(CVC) + P(CCV) + P(CCC)$

$$\begin{aligned} &= \frac{3}{8} \times \frac{5}{8} \times \frac{4}{7} + \frac{5}{8} \times \frac{3}{7} \times \frac{4}{7} + \frac{5}{8} \times \frac{4}{7} \times \frac{3}{6} + \frac{5}{8} \times \frac{4}{7} \times \frac{3}{6} \\ &= \frac{505}{784} \\ &= 0.644 \end{aligned}$$

9(i)

Using GC,

Unbiased estimate of population mean is $\bar{x} = 949.84$ (2 d.p.)

Unbiased estimate of population variance is $s^2 = 1.31634^2 = 1.73$ (2 d.p.)

Note: If student write $\mu = 949.84$ or

$\sigma^2 = 1.31634^2 = 1.73$, annotate but do not deduct.

M
di

G
St
ar
gi
T
m
st
1.
gc

2.

<p>9 (a) (ii)</p>	<p>Let X be the volume of shower gel dispensed by the machine.</p> <p>Let μ denote the population mean volume of shower gel dispensed by the machine.</p> <p>$\bar{X} \sim N\left(\mu, \frac{s^2}{n}\right)$ approx. by Central Limit Theorem.</p> <p>$\bar{X} \sim N\left(950, \frac{s^2}{n}\right)$</p> <p>$H_0: \mu = 950$</p> <p>KIASU ExamPaper950 Islandwide Delivery Whatsapp Only 88660031</p>	<p>N 1. σ sa U va (1 (1 fo N μ Pa ar M th de 's de of ex th as er</p>
<p>9 (a)(iii)</p>	<p>Under H_0, using GC,</p> <p>p-value = 0.13474 \approx 0.135</p>	<p>R m fa st cc</p>

	<p>The p - value is the probability of the sample mean volume of shower gel in a bottle is as extreme as 949.84 when the mean is actually 950.</p>	<p>Th w 9 — m M th cc So th w of hy st p- cc</p>
<p>9 (a)(iv)</p>	<p>Since we reject H_0 , p - value $< \alpha / 100$</p> <p>$\alpha \geq 13.476$ $\{\alpha \in \mathbb{R} : 13.5 \leq \alpha \leq 100\}$</p>	<p>B di nc up St us in ro th</p>
<p>9(b)</p>	<p>Let Y be the volume of shower gel dispensed by the machine after recalibration.</p> <p>Let μ denote the population mean volume of refill dispensed by the machine.</p> <p>$\bar{Y} \sim N\left(\mu, \frac{s^2}{n}\right)$ approx. by Central Limit Theorem</p> <p>$H_0: \mu = 250$ $H_1: \mu \neq 250$</p> <p>Test Statistic: $\frac{\bar{Y} - \mu}{s/\sqrt{n}}$</p> <p>Level of significance: 1%</p> <p>Reject H_0 if p - value ≤ 0.01</p>	<p>C th 1. 2. to 3. as cc 4.</p>

	$\frac{249.5 - 250}{s/\sqrt{50}} \leq -2.5758293 \text{ or } \frac{249.5 - 250}{s/\sqrt{50}} \geq 2.5758293 \text{ (NA)}$ $0 < \frac{s}{\sqrt{50}} \leq 0.19411$ $0 < s \leq 1.3726$ $0 < \frac{1}{49} \left(k - \frac{(-25)^2}{50} \right) \leq 1.3726^2$ $12.5 < k \leq 104.8$	M lo w in M th 2 — re th																									
9(c)	<p>There is no need for the floor supervisor to assume the volume of shower gel follow a normal distribution as the sample sizes in both part (a) and (b) are large. The sample mean volume of shower gel can be approximated to follow a normal distribution by Central Limit Theorem.</p>	A Tl sa ca no Tl w 's St ar fo																									
10 (i)	<p>Let \$W\$ denote the amount won by a player in one game. Under option A,</p> $E(W_A) = 1\left(\frac{1}{4}\right) + 2\left(\frac{1}{4}\right) + 3\left(\frac{1}{4}\right) + 4\left(\frac{1}{4}\right)$ $= (1+2+3+4)\left(\frac{1}{4}\right)$ $= \frac{5}{2} = \$2.50$ <p>Under option B, the winning amount for each combination is listed below:</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>1</th> <th>2</th> <th>3</th> <th>4</th> </tr> </thead> <tbody> <tr> <th>1</th> <td>0</td> <td>4</td> <td>6</td> <td>8</td> </tr> <tr> <th>2</th> <td>-1</td> <td>0</td> <td>12</td> <td>16</td> </tr> <tr> <th>3</th> <td>-2</td> <td>-1</td> <td>0</td> <td>24</td> </tr> <tr> <th>4</th> <td>-3</td> <td>-2</td> <td>-1</td> <td>0</td> </tr> </tbody> </table>		1	2	3	4	1	0	4	6	8	2	-1	0	12	16	3	-2	-1	0	24	4	-3	-2	-1	0	St ra di ne ar pr sc G ex A H nu kr w re as po di cc
	1	2	3	4																							
1	0	4	6	8																							
2	-1	0	12	16																							
3	-2	-1	0	24																							
4	-3	-2	-1	0																							

	$E(W_B) = \left(\frac{1}{16}\right)[4+6+12+8+16+24$ $-1-2-3-1-2-1]$ $= \left(\frac{1}{16}\right)[70-10]$ $= \frac{15}{4} = \$3.75$ <p>Since $E(W_B) > E(W_A)$, option B is better. (shown)</p>	ca ar It ex ar O
(ii)	Option A is a “sure win” option where the player would definitely gain a positive amount in all cases, whereas option B has a risk of losing money in some cases.	G
(iii)	$E(W_A^2) = (1^2 + 2^2 + 3^2 + 4^2) \left(\frac{1}{4}\right) = \frac{15}{2}$ $\text{Var}(W_A) = E(W_A^2) - [E(W_A)]^2 = \frac{15}{2} - \left(\frac{5}{2}\right)^2 = \frac{5}{4} = 1.25 \text{ (shown)}$	G Tl st cc
(iv)	<p>Let A and B denote the total amount won by Abel and Benson respectively.</p> $A = W_{A1} + W_{A2} + \dots + W_{A50}$ <p>Since $n = 50$ is large, by Central Limit Theorem,</p> $A \sim N\left(50\left(\frac{5}{2}\right), 50\left(\frac{5}{4}\right)\right) \text{ approximately}$ <p>i.e. $A \sim N\left(125, \frac{125}{2}\right)$</p> $B = W_{B1} + W_{B2} + \dots + W_{B50}$ $B \sim N\left(50\left(\frac{15}{4}\right), 50\left(\frac{887}{16}\right)\right) \text{ approximately by Central}$ <p>Limit Theorem since $n = 50$ is large</p> $B \sim N\left(375, \frac{22175}{8}\right)$	A w va ar w th Tl ap of ar O - - - -

		-
(v)	$A - B \sim N\left(125 - \frac{375}{2}, \frac{125}{2} + \frac{22175}{8}\right)$ $A - B \sim N\left(-\frac{125}{2}, \frac{22675}{8}\right)$ $P(A > B) = P(A - B > 0)$ $= 0.120207$ $= 0.120(3sf)$	M at re pr A cc ar fo - -